

NEW
SOFT BAIT



CONTRAC[®]

SOFT BAIT

NET WT:
16 lbs (7.2 kg)

KILLS RATS, MICE & MEADOW VOLES*
Kills Warfarin Resistant Norway Rats And House Mice

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for First Aid and additional precautionary statements.

ACTIVE INGREDIENT:	
Bromadiolone (CAS #28772-51-7):	0.005%
OTHER INGREDIENTS:	99.995%
*Contains Denatonium Benzoate	TOTAL 100.000%

*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat

FIRST AID
HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE

IF SWALLOWED:
 • Have person sip a glass of water if able to swallow.
 • Do not induce vomiting unless told to do so by the poison control center or doctor.

IF ON SKIN OR CLOTHING:
 • Take all contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
 • If in eyes:
 • Hold eye open and flush slowly and gently with water for 15-20 minutes.
 • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Call a poison control center, doctor, or 1-877-454-2486 immediately for treatment advice.

NOTE TO PHYSICIAN
 If swallowed or absorbed through the skin, this bait may reduce the clotting ability of the blood and cause bleeding. If repeated, administer Vitamin K1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin time.

TREATMENT FOR PET POSIONING
 If animal eats bait, call veterinarian at once.

NOTE TO VETERINARIAN
 Anticoagulant Bromadiolone: For animals exhibiting bait and/or showing poisoning signs (bleeding or abnormal prothrombin time), give Vitamin K1. If needed, check prothrombin time every 2 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
READ THIS LABEL: Read the entire label and follow all use directions and use procedures. Use only for the uses, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets or nontarget animals to rodenticides. To help to prevent exposure:

1. Store unused product out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. Bait can be obtained from bait stations when they are food units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hooded rhesus monkeys, bears or other potentially destructive animals or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled, or contaminated bait as specified on this label.

Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used whenever children, pets, nontarget mammals, or birds may have access to the bait placement location.

USE RESTRICTIONS: This product may only be used to control the following rodent pests: *House mouse* (*Mus musculus*), *Norway rat* (*Rattus norvegicus*), *Roof rat* (*Rattus rattus*), *Common rat* (*Sigmodon hispidus*), *Eastern harvest mouse* (*Reithrodontomys fulvus*), *Golden mouse* (*Ochrotomys nuttalli*), *Polynesian rat* (*Rattus exulans*), *Norfolk vole* (*Microtus pennsylvanicus*), *White-throated woodrat* (*Microtus agilis*), *Southern plains woodrat* (*Neotoma albigula*), and *Hispanian woodrat* (*Neotoma mexicana*). This product must be used in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commercial rodent infestations and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other residences, or temporary residences, food processing facilities, industrial and commercial buildings, retail establishments, agricultural and public buildings, transport vehicles (trucks, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops.

577CS



KILLS RATS, MICE & MEADOW VOLES
Kills Warfarin Resistant Norway Rats and House Mice

Honey rats, roof rats, and house mice may consume a lethal dose in one night feeding with first bait contact, appearing four or five days after feeding begins.

ACTIVE INGREDIENT: Bromadiolone (CAS #28772-56-7)	0.005%
OTHER INGREDIENTS:	89.995%
1 Contains Denaletum Benzothiole	TOTAL 100.000%

KEEP OUT OF REACH OF CHILDREN
CAUTION
 Use side panel for First Aid and additional precautionary statements.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Plastic: Storage: Store only in original container and seal tightly place inaccessible to children and pets. Keep container closed and away from open flames.
Pesticide Disposal: Waste resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.
Container Handling: Reusable container. Do not reuse or refill this container. [Plastic:] Offer for recycling or reconditioning or purchase and dispose of in a primary landfill, or by incineration. In most states, burning is not allowed. [Paper:] Dispose of empty container by placing in trash at an approved waste disposal facility or by incineration. In most states, burning is not allowed.

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat



NET WEIGHT: 16 lbs (7.2 kg)
 0.35 oz (10g) sachets

EPA REG. NO. 12455-W-0
 EPA EST. NO. 12455-W-11
 12455-W-02P, 12455-W-03P
 Supplemental is trial letter of the lot number.

MADE IN USA

USE RESTRICTIONS: (Continue from left page)
 This product may also be used in alfalfa. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feed, or food handling equipment or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations. Do not broadcast bait.
 Do not sell this product in individual containers holding less than 16 pounds of bait.
 Baiting with Contract® Soft Bait is prohibited.
SELECTION OF TREATMENT AREAS: Determine areas where use, mice, or meadow voles will most likely feed and consume the bait. Generally, these areas are along walls, by ground openings, in corners and concealed places, between floor and walls, inside burrows, or in locations where rats, mice, or meadow voles or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.
APPLICATION DIRECTIONS:
RATS: Place 9 to 43 sachets per placement usually at intervals of 15 to 30 feet in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.
MICE AND MEADOW VOLES: Place 1 to 2 sachets per placement usually at intervals of 8 to 12 feet in infested areas. Up to 5 sachets may be needed at points of very high activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse or meadow vole activity cease.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Wearing gloves collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur repeat treatment. While a continuous source of infestation is present, establish permanent bait stations and maintain as needed.

Bait may be removed from sachet and placed in a feeding device that secures the unwrapped bait in a bait station. Discard empty wrapper in the trash.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed or absorbed through the skin. Keep away from children, domestic animals and pets. Do not get in eyes, on skin or on clothing.
 All handlers (including applicators) must wear shoes plus socks and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

User Safety Requirements
 Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

ENVIRONMENTAL HAZARDS
 This product is extremely toxic to fish, birds and other wildlife. Dogs, cats, and predatory scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply this product directly to water or to areas where surface water is present or to littoral areas below the mean high water mark. Runoff also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or residue.

050317/02-17

Product Code: CS1016