

InTice™

sweet ant gel

KILLS SWEET FEEDING ANTS

Doesn't Drip • Stays Soft

Active Ingredient: Orthoboric Acid.....5%
Inert Ingredients:.....95%
Total:.....100%

EPA Reg. No. 73079-1
EPA Est. No. 73079-MO-1
Net Contents: 5 x 1.5 oz Syringes

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

- IF SWALLOWED:**
- Call a poison control center or doctor immediately for treatment advice.
 - Have a person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by a poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.
- IF IN EYES:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.
- IF ON SKIN OR CLOTHING:**
- Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.

Have the product or label with you when calling a Poison Control Center or doctor or going for treatment. In case of emergency, for additional information call **800 858 7378**.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION Harmful if swallowed. May cause eye irritation. Avoid contact with skin, eyes, or clothing. Wash hands thoroughly after handling.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Sweet feeding ants are usually found in moist areas such as kitchens and bathrooms, and will often appear to be randomly searching for food. Baits will work best if competing food sources are cleaned up and placed in sealed containers. This product is designed for use on sweet feeding ants. Place bait in areas inaccessible to children and pets or in tamper-resistant bait stations.

To locate ant trails: First locate foraging ants and place a drop of InTice Sweet Ant Gel in front of an ant. When the ant makes contact, elongate the drop to 1/2 inch. Normally within 10-20 minutes, the ants will recruit more workers, forming a trail that can be followed to locate where ants are coming into the structure. Check other areas of the structure to determine if there are other entry points, potentially indicating more than one nest.

Baiting: Bait ants close to their entry point(s) and along their trail(s). Long-term success will come from baiting each separate ant trail. To maximize feeding surface, apply bait in thin (1/8 inch wide) lines, one to two inches long. Use one to three lines per entry point or trail. After baiting, lightly blow on ants feeding at the pre-treat spot to disperse, and wipe up the bait with warm water. Replenish bait as needed. Clean up any remaining bait with warm water at the end of the baiting program. Results are enhanced if the same procedure is used to bait around the outside of the structure.

The non-drip formula of InTice Sweet Ant Gel makes it easy to bait out of the way on vertical and horizontal surfaces such as the edges and tops of cupboards, along window frames and in corners where the wall and floor or ceiling meet. Some water marking may occur on unfinished surfaces. In the edible product areas of food handling establishments, other areas where food is commercially prepared or processed, and in food service areas, only place bait in cracks and crevices or covered bait stations and record the number and location of bait stations. Avoid contamination of feed and foodstuffs.

STORAGE AND DISPOSAL

STORAGE: Store in a dry place inaccessible to children and pets.

DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP which is managed as a public-private partnership.

MANUFACTURED BY:



1512 Taney St. • North Kansas City, MO 64116
toll free: 866 788 4101 • fax: 816 283 3173
www.rockwelllabs.com